

U.S. DEPARTMENT OF COMMERCE Patent and Trademark Office

Complete if Known Application Number 10/612,468 Filing Date 07/02/2003 First Named Inventor Jingwu Z. Zhang et al. Group Art Unit Unknown Examiner Name Unknown

INFORMATION DISCLOSURE STATEMENT BY APPLICANT

 (use as many sheets as necessary)
 Attorney Docket Number
 057186.000003

U.S. PATENT DOCUMENTS

Examiner Initials	Cite No.	U.S. Pate	ent Document	Publication Date MM-YYYY	Name of Patentee or Applicant of Cited Document	Pages, Columns, Lines, where Relevant Passages or Relevant Figures Appear
		Number	Kind Code (if known)			

				FORE	EIGN PATENT DO	CUMENTS	
Examiner Initials	Cite No.	F	oreign Patent Docum	nent	Publication Date	Name of Patentee or Applicant of Cited Document	Pages, Columns, Lines Where Relevant Passages or Relevant Figures Appear
U N U GI J	140.	Country Code	Number	Kind Code (if known)			•
		l l		1		!	

	· ·	OTHER PRIOR ART
Examiner nitials	Cite No.	Include Name of Author (in CAPITAL LETTERS), title of the article (when appropriate), title of the item (book, magazine, journal, serial, symposium, catalog, etc.), date, page(s), volume-issue number(s), publisher, city and/or county where published.
AJ		Lee DM, Weinblatt ME. Rheumatoid arthritis. Lancet 2001;15:903-911.
		Firestein GS. Evolving concepts of rheumatoid arthritis. Nature 2003;423:356-361.
		Kerian-Candon S, Combe B, Vincent R, Clot J, Pinet V. Eliaou JF. HLA-DRB1 gene transcripts in
		rheumatoid arthritis. Clin Exp Immunol 2001;124:142-149.
		MacGregor A, Oilier W, Thomson W, Jawaheer D, Silman A. HLA~DRBI*0401/0404 genotype
		and rheumatoid arthritis: increased association in men, young age at onset, and disease severity. J
		Rheumatol 1995;22:1032-1036.
		Fries JF, Wolfe F, Apple R, Erlich H, Bugawan T, Holmes T, Bruce B. HLA-DRB1 genotype
		associations in 793 white patients from a rheumatoid arthritis inception cohort: frequency, severity
- 1		and treatment bias. Arthritis Rheum 2002;46:2320-2329.
		Dolhain RJEM, van der Heiden AN, Ter haar NT, Breedveld FC, Miltenburg AM. Shift towards T
1		lymphocytes with a T helper 1 cytokine profile in the joints of patients with rheumatoid arthritis.
		Arthritis Rheum 1996; 12:1961-1969.
		Bemer B, Akca D, Jung T, Muller GA, Reuss-Borst MA. Analysis of Thi and Th2 cytokines
	ŀ	expressing CD4+ and CD8+ T cells in rheumatoid arthritis by flow cytometry J Rheumatol
		2000;27:1 128-1135.
		Davis LS, Cush JJ, Schultz-Koops H, Lipsky PE. Rheumatoid synovial CD4+ T cells exhibit a
		reduced capacity to differentiate into IL-4 poducing T helper-2 effector cells. Arthritis Res
		2001;3:54-64.
АJ		Goronzy JJ, Bartz-Bazzanella P, Hu W, Jendro MC, Walser-Kuntz DR, Weyand CM. Dominant
		clonotypes in the repertoire of peripheral CD4+ T cells in rheumatoid arthritis, J Clin Invest
		1994;94:2068-2076.

i da a	Goronzy JJ, Bartz-Bazzanella P, Hu W, Jendro MC, Walser-Kuntz DR, Weyand CM. Dominant
ADEMEAJ	clonotypes in the repertoire of peripheral CD4+ T cells in rheumatoid arthritis. J Clin Invest 1994;94:2068-2076.
	Gonzalez-Quintial R, Baccala R, Pope RM, Theofilopoulos AN. Identification of clonally expanded T cells in rheumatoid arthritis using a sequence enrichment nuclease assay. J Clin Invest 1996;97:1335-1343.
	Alam A, Lambert N, Lule H, Coppin H, Mazieres B, de Preval C, et al. Persistence of dominant T cell clones in synovial tissues during rheumatoid arthritis. J Immunol 1996; 156:3480-3485.
	Londei M, Savill CM, Verhoef A, Brennan F, Leech ZA, Duance V, et al. Persistence of collagen type IT-specific T-cell clones in the synovial membrane of a patient with rheumatoid arthritis. Proc Nati Acad Sci USA 1989;86:636-640.
	Pope RM, Pahiavani MA, LaCour E, Sambol S, Desai BY. Antigen specificity of rheumatoid synovial fluid lymphocytes. Arthritis Rheum 1 989;32:1371-1380.
	Devereux D, O'Hehir RE, McGuire J, van Schooten WC, Lamb JR. HLA-DR4Dw4-restricted T cerecognition of self antigen(s) in the rheumatoid synovial compartment. Int Immunol 1991;3:635-640.
	Res PC, Struijk L, Leow A, Daha MR, van den Elsen PC, Breedveld FC. Inflamed joints of patients with rheumatoid arthritis contain T cells that display in vitro proliferation to antigens present in autologous synovial fluid. Functional analysis on the basis of synovial-fluid-reactive T cell clones and lines. Hum Immunol 1994;40:291-298.
	Paliard X, West SF, Lafferty JA, Clements IR, Kappler JW, Marrack P, et al. Evidence for the effects of a superantigen in rheumatoid arthritis. Science 1991;253:325-329.
	Holoshitz J, Klajman A, Drucker I, Lapidot Z, Yaretzky A, Frenkel A, et al. T lymphocytes of rheumatoid arthritis patients show augmented reactivity to a fraction of mycobacteria cross-reactive with cartilage. Lancet 1986;2:305-309.
	Zagon G, Tumang JR, Li Y, Friedman SM, Crow MK. Increased frequency of V beta 17-positive I cells in patients with rheumatoid arthritis. Arthritis Rheum 1994;37:1431-1440.
	Alam A, Lule J, Coppin H, Lambert N, Mazieres B, De Preval C, et al. T -cell receptor variable region of the beta-chain gene use in peripheral blood and multiple synovial membranes during rheumatoid arthritis. Hum Immunol 1995;42:331-339.
	VanderBorght A, Geusens P, Vandevyver C, Raus J, Stinissen P. Skewed T-cell receptor variable gene usage in the synovium of early and chronic rheumatoid arthritis patients and persistence of clonally expanded T cells in a chronic patient. Rheumatology 2000;39:1 189-1201.
	Jenkins RN, Nikaein A, Zimmermann A, Meek K, Lipsky PE. T cell receptor V beta gene bias in rheumatoid arthritis. J Clin Invest 1993;92:2688-2701.
	Williams WV, Fang Q, Demarco D, VonFeldt J, Zurier RB, Weiner DB. Restricted heterogeneity of T cell receptor transcripts in rheumatoid synovium. J Clin Invest 1992;90:326-333.
	Li Y, Sun GR, Tumang JR, Crow MK, Friedman SM. CDR3 sequence motifs shared by oligoclona rheumatoid arthritis synovial T cells. Evidence for an antigen-driven response. J lin Invest 1994;94:2525-2531.
	Mima T, Ohshima 5, Sasi M, Nishioka K, Shimizu M, Murata N, et al. Dominant & shared T cell receptor beta chain variable regions of I cells inducing synovial hyperplasia in rheumatoid arthritis. Biochem Biophys Res Commun 1999;263:172-180.
	Davey MP, Burgoine GA, Woody CN. TCRB clonotypes are present in CD4+ T cell populations prepared directly from rheumatoid arthritis. Hum Immunol 1 997;55: 11-21.
	Even J, Lim A, Puisieux I, Ferradini L, Dietrich PY, Toubert A, et al. T-cell repertoires in healthy and diseased human tissues analysed by T-ce!! receptor beta-chain CDR3 size determination: evidence for oligoclonal expansions in tumors and inflammatory diseases. Res Immunol 1995; 146:65-80.
\downarrow	Tang J, Myracle AD, Allen S, Karita E, Musonda R, Fultz PN, et al. Novel alleles at the lymphotoxin alpha (LTalpha) locus mark extended HLA haplotypes in native Africans. Hum Immunol 2001;62:269-278.
AJ	Nakao M, Janssen JW, Flohr T, Bartram CR. Rapid and reliable quantification of minima! residual disease in acute lymphoblastic leukemia using rearranged immunoglobulin and I-cell receptor loci by LightCyc!er technology. Cancer Res. 2000;60:3281-289.

O 1 P 4	****	
DCT 14	AJ AJ	Hong J, Zang YC, Tejada-Simon MV, Kozovska M, Li S, Singh RA, et a!. A common TCR V-D-J sequence in V beta 13.1 T cells recognizing an immunodominant peptide of mye!in basic protein in multiple sclerosis. J Immunol 1999;163:3530-3538.
VA	AJ	Moreland LW, Morgan EE, Adamson TC 3rd, Fronek Z, Calabrese LH, Cash JM, et al. T cell receptor peptide vaccination in rheumatoid arthritis: a placebo-controlled trial using a combination of Vbeta3, Vbeta14, and Vbeta17 peptides. Arthritis Rheum 1998;41:1919-1929.
	AJ	Moreland LW, Heck LW Jr, Koopman WJ, Saway PA, Adamson IC, Fronek Z, et al. V beta 17 T cell receptor peptide vaccination in rheumatoid arthritis: results of phase I dose escalation study. J Rheumatol 1996;23:1353-1362.

Signature / 12mly Steedes/	Examiner Signature /Amy Juedes/ Date considered 05/
----------------------------	---